

Reorganization of columnar architecture in the growing visual cortex

Wolfgang Keil,^{1,*} Karl-Friedrich Schmidt,² Siegrid Löwel,² and Matthias Kaschube^{3,†}

¹*Max-Planck Institute for Dynamics and Self-Organization, 37073 Göttingen, Germany*

²*Institute of General Zoology and Animal Physiology,
Friedrich-Schiller University, 07743 Jena, Germany*

³*Lewis Sigler Institute for Integrative Genomics, Princeton University, 08544 Princeton, NJ*

Many cortical areas increase considerably in size during postnatal development, progressively displacing neuronal cell bodies from each other. At present it is unclear to what extent this requires remodeling of neuronal circuits. Here, in acute and chronic experiments, we study the layout of ocular dominance columns (OD) in cat primary visual cortex (V1) during a period of substantial postnatal growth. We find that OD columns reorganize such that the spacing between columns is largely unaffected, despite a considerable size increase of V1. We show that the observed mode of reorganization can be explained by a fundamental mechanism of growth induced reorganization inherent to a very general class of models of activity-dependent formation of OD columns. Our work suggests that cortical circuits remain plastic for an extended period in development in order to facilitate the modification of neuronal circuits to adjust for cortical growth.

I. INTRODUCTION

The brain of most mammalian species grows substantially during postnatal development. The human brain, for instance, weighs on average 350 g in newborns and 1400 g in adult males [1]. In cat, the neocortical volume increases from $\approx 1000 \text{ mm}^3$ at birth to $\approx 4500 \text{ mm}^3$ in adulthood [2]. Consistently, the surface area of cat primary visual cortex (V1) increases postnatally by 130% between week 1 and week 15 [3]. The number of neurons remains largely constant during this period [4]. Most of the area increase is due to the generation of glial cells, the addition of more vasculature and connective tissue, and the myelination of axons; to a lesser extent it also reflects the outgrowth and elaboration of axonal and dendritic processes [5].

It is likely that such substantial cortical growth induces remodeling of neuronal circuits. The size increase of cat V1 implies that the distance between any two neuronal cell bodies grows on average by $\approx 50\%$ during early postnatal development. Consequently, synaptic connections between cortical neurons that are established early in development must be either replaced, or dendritic and axonal processes must grow by a similar factor in order to preserve these connections. However, such morphological changes, albeit subtle, can modify the temporal properties of neurons. For instance, significant dendritic growth of a pyramidal neuron in cat visual cortex could induce delays in dendritic integration [6] on the order of a millisecond assuming a lateral dendritic spread of $\approx 300 \mu\text{m}$ [7] and propagation speeds of voltage changes on the order of $100 \mu\text{m/ms}$ [8]. Moreover, since synaptic plasticity can be sensitive on a millisecond time scale [9], this might have long lasting effects on synaptic efficacy. These growth-induced changes potentially apply

to a large fraction of synaptic connections and thus may affect significantly patterns of cortical activity.

Much of the growth of V1 takes place during a period when many neurons have already reached fairly mature levels of selectivity. For instance, the selective response of visual cortical neurons to inputs from one eye or the other, called ocular dominance (OD), can already be visualized at postnatal week 2 in cat V1 [10]. OD is organized into columns which can be labeled within the entire V1 as early as week 3 [11, 12]. Large parts of the visual field are already represented at this time [13]. Subsequently, between week 3 and week 10, cat V1 increases its size by 50% [12, 14], a result which we confirm in **Figure 1**. In this work, we exploit the fact that OD columns can be visualized over a period of significant cortical growth to reveal signatures of growth induced remodeling of neuronal circuits.

We analyze the layout of OD columns during cortical growth in experiment and in models for the activity dependent formation of OD columns. Complete reconstructions of OD columns in cat V1 are obtained using 2-[¹⁴C]-deoxyglucose (2-DG) and [³H]-proline labelling of OD columns. Additionally, we follow the development of columnar layouts in individual animals by performing chronic optical imaging experiments. We find that, long after OD columns have segregated, columnar layouts reorganize throughout V1. Despite the considerable area increase of V1, the spacing of OD columns is largely preserved. To understand these empirical observations, we study growth induced reorganization theoretically in models for activity-dependent formation of OD columns. We argue that in a very large class of models cortical growth induces a characteristic zigzag-like bending of OD columns by which their spacing is largely restored. This class contains, to our best knowledge, any model that has been proposed thus far (e.g. [15, 16, 17, 18, 19]). As a representative of this class, we study in detail the reorganization of OD layouts in different growth scenarios in the Elastic Network model [19, 20]. We find that for wide range of growth rates, the growth induced reorga-

*Electronic address: wolfgang@nld.ds.mpg.de

†Electronic address: kaschube@princeton.edu

nization in the model shows the same characteristics as the reorganization observed in experiment. Our results suggest that cortical growth accompanied by significant displacement of neuronal cell bodies can induce substantial remodeling of neuronal circuits. Furthermore, we suggest a possible new role of the critical period in normal development which in cat visual cortex overlaps with the period of postnatal growth (**Fig. 1c**): The enhanced degree of plasticity during the critical period may facilitate the modifications of neuronal circuits to adjust for cortical growth.

II. RESULTS

OD Column Spacing Changes Only Little During Postnatal Growth of Primary Visual Cortex

We first measured the size increase of cat primary visual cortex (V1) during early postnatal development. We labelled complete layouts of OD columns in V1 visualized by either 2-DG or proline in kittens at different ages between postnatal day (PD) 30 and PD84 ($N = 18$). The left panel of **Figure 1a** displays a representative 2-DG autoradiograph of a kitten at PD30. The right panel displays proline labeled OD columns of a kitten at PD60. The range of our entire data set compared with the time line of cat postnatal development is displayed in **Figure 1b**. V1 was readily discernable by its distinctive columnar activation pattern in comparison to the labelling in surrounding cortical areas (see also [21, 22]). We observed a size increase of about 30% between two groups centered at PD30 and PD70 ($r = 0.6248$, $p < 0.006$) (**Fig. 1b**). To minimize possible influences of genetic variability [23], we analyzed a littermate couple on PD30 and PD72 respectively (green crosses in **Fig. 1b**). Consistently, their cortical sizes also differed by an even larger factor of 1.46. Hence, our analysis confirmed the considerable size increase of V1 during early postnatal development observed in previous studies [3, 12].

Given this size increase of V1 during the first postnatal weeks, we next asked whether the spacing of OD columns increases by a corresponding factor over this time period. Such a finding would be consistent with a scenario that involves no change of column number or layout (sometimes referred to as “balloon effect” [12, 24]). First, we measured the mean column spacing Λ of 2-DG/proline labelled OD patterns in $N = 41$ hemispheres (2-DG, $N = 37$; proline, $N = 4$) between PD28 and PD98 (data includes the $N = 18$ hemispheres used for the analysis of V1 sizes). To obtain accurate estimations of column spacings, we used a wavelet-method that exhibits a precision of 2% (relative error, see [23, 25] and **Methods**). As shown in **Figure 2a**, mean column spacings varied between 1.05 mm and 1.28 mm, but did not show a significant increase over this period ($r = -0.034$, $p < 0.825$). Consistent with this general trend, the column spacings of the two littermates differed by less than 10%, despite

a size difference of almost 50%.

To follow the development of OD column spacing in individual animals we visualized OD columns in cat V1 by chronic optical imaging experiments over the same period ($N = 3$ hemispheres, total range: PD30-PD94). **Figure 2b** shows the layout of OD columns at different ages in one of the studied cases (data are high-pass filtered, see **Methods**). Again, we quantified column spacings of OD columns by the wavelet-method. As shown in **Figure 2c**, we found no systematic increase of column spacings in individual animals, thus confirming the conclusions drawn from the 2-DG/proline data. Column spacing estimates based on optical recordings exhibited a larger variability compared to the 2-DG/proline data. This might be explained by the substantial interareal variability of OD column spacing [25] and the fact that the recorded regions spanned only a small portion of V1 and possibly shifted with age. Taken together, both the 2-DG/proline data and the chronic optical recordings demonstrate that the postnatal growth of the primary visual cortex is not accompanied by a corresponding increase in the spacing of OD columns.

A Universal Mode of Reorganization

At first glance, the finding of postnatal cortical growth without a corresponding increase in OD column spacing might seem contradictory. However, we will show in the following sections that both findings can naturally be reconciled within a very large class of models for activity-dependent formation of OD columns.

In most models (e.g. [15, 16, 17, 18, 19]), the segregation of OD columns is caused by a Turing mechanism [26], also called finite wavelength instability [27]. Typically, the Turing mechanism results from Mexican-Hat-like cortical interactions (whether assumed explicitly or arising effectively from the assumed dynamics). By this mechanism, a layer of nonselective neurons segregates into alternating domains of roughly similar size selective for one eye or the other. The spacing between these domains is approximately twice the distance between the maximum and the minimum of the Mexican-Hat.

How does cortical growth affect columns that already have segregated, according to these models? One possibility would be that the size of the effective Mexican-Hat scales with the size of the cortex and increases by the same factor. This could keep neuronal interactions unchanged, but would also increase column spacing by the same factor. Since we found no evidence for such an increase of column spacing, we conclude that this scenario is rather unlikely.

Suppose that instead the effective Mexican-Hat does not increase or increases less than the cortical tissue. Here we conjecture that in this case a reorganization of OD domains takes place that shows signatures of the so called zigzag (ZZ) instability. The ZZ instability has been observed in many inanimate dynamical pattern form-




Figure 1: Postnatal growth of cat primary visual cortex (V1). (a) Two representative examples of OD layouts in kittens with different age. Left/right panel shows a 2-DG/proline autoradiograph of kittens at PD30/PD60. Yellow lines mark V1 borders. Scale bar, 10 mm. (b) Area sizes of V1 in kittens between PD30 and PD84. The average size increases by $\approx 30\%$ percent between the two groups centered around PD30 and PD70. Red line, linear fit ($r = 0.6248$, $p < 0.006$). The two green crosses mark values from a littermate couple. (c) Growth of cat V1 with representation of the visual hemi-field (schematic; HM, horizontal meridian; VM, vertical meridian) and time line of cat OD development. OD segregation can be visualized as early as PD15 [10], OD critical period for monocular deprivation starts at PD21 [10]. Our dataset includes 2-DG/proline labelled hemispheres from kittens between PD28 and PD98 and chronic optical recordings between PD30 and PD96




Figure 2: Spacings of OD columns change only little during growth of cat visual cortex. (a) Mean column spacing Δ of OD columns for $N = 41$ hemispheres of cat V1 between PD28 and PD98 (2-DG, $N = 37$; proline, $N = 4$). The two green crosses mark values from two littermates. Red line, linear fit ($r = -0.034$, $p < 0.825$). (b) Chronic recordings of OD columns in an individual hemisphere at PD30, PD42 and PD94 (intrinsic signal optical imaging, $N = 3$, scale bar 1.5 mm). Note that the recorded region is only 4-5 column spacings wide and may have shifted during development. (c) Mean column spacings for the case in b (blue circles) and for two other cases (red boxes and green crosses). Column spacings from a are shown in light gray for comparison.

ing systems as, for instance, Rayleigh-Benard convection, when these systems experience an instantaneous isotropic size increase [27]. The result of a ZZ instability can be best understood by considering a simple pattern of OD columns consisting of alternating stripes

(Fig. 3a). After an isotropic area increase, the spacing between the OD stripes is larger than the spacing set by the Mexican-Hat. Subsequently, the structure rearranges in a sinusoidal fashion: A zigzag-like bending occurs by which the original spacing is largely recovered, locally.

Figure 3b illustrates this in Fourier space. The initial stripe pattern is described by a wave $\sim e^{ik_{\max}\tilde{\mathbf{x}}}$ with spatial wavenumber $k_{\max} = 2\pi/\Lambda_{\max}$. Upon area increase, its wave vector differs by an amount $\delta k < 0$ from $k_{\max}$, i.e. the spacing is larger than $\Lambda_{\max}$. Two new Fourier modes grow with wave vector $(k_{\max} + \delta k)\tilde{\mathbf{x}} \pm q_y\tilde{\mathbf{y}}$ representing the parts of the OD stripes bended to the left and right, respectively (**Fig. 3a**). The diagram in **Figure 3c** shows the region of the ZZ instability as a red shaded region in the r - δk plane, where r is the control parameter of the system expressing the distance from the primary (Turing) instability (for a more detailed description, see **Methods**). The diagram also shows that insertion of a new stripe or elimination of an existing stripe requires strong area increase or decrease, respectively (the regions of Eckhaus (EH) instability). The form of this diagram is quite general. It is typical of all rotationally invariant systems that exhibit a Turing instability and it does not reflect the details of the underlying microscopic equations [27].

Thus, for all models of OD formation that are rotationally symmetric and based on a Turing mechanism, we expect to observe a reorganization of columnar layouts upon isotropic area increase that shows signatures of ZZ instability. Most models that have been proposed (e.g. [15, 16, 17, 18, 19]) fall into this class. In the following, we focus on one of its representatives, namely the Elastic Network (EN) model [19, 20, 28]. This model has been widely used to describe the formation of visual cortical selectivity patterns and here serves as a concrete example for studying in detail growth induced reorganization.

Since in a realistic growth scenario the cortex increases continuously in size and the initial OD layout is not a simple stripe pattern, the growth induced reorganization may be more complex than an ideal ZZ instability. However, we expect that the reorganization of OD layouts still shows signatures of a ZZ instability such as sinusoidal bending or distortion of local pattern elements. For simplicity, we refer to all growth induced changes as ZZ reorganization in the following. To capture quantitatively changes in local map layout both in model and experiment we calculate a parameter called bandedness α [23] that quantifies the tendency of local pattern elements to be composed of parallel stripes (see **Methods** and **Fig. 10**). Large values of α indicate layouts composed of regular stripe like parallel domains, whereas small values indicate bended, non-parallel stripes or patches. Together with the column spacing, Λ , and the bandedness α , we use the number of hypercolumns $N_{HC} = A/\Lambda^2$ in an area of size A [25]. The quantity Λ^2 is the size of the typical unit cell of an OD map, which we refer to as the hypercolumn size. Before studying more realistic growth scenarios, we consider a near stripe like OD steady state solution for the EN model subject to instantaneous area increase and reproduce numerically the stability diagram sketched in **Figure 3c**. First, however, we study the EN model without assuming any growth.




Figure 3: What do models predict about OD development in a growing visual cortex? Assuming that cortical interactions do not scale with growth (**Fig. 2a, c**), most models [15, 16, 17, 18, 19] predict a zigzag (ZZ) reorganization that induces sinusoidal bending of domains (schematic). This is a simple mechanism by which the OD stripes can recover their column spacing Λ_1 . (a) During isotropic area increase, an OD stripe pattern transiently acquires a spacing $\Lambda_2 = \Lambda_1 + \delta\Lambda$, larger than Λ_1 . By the ZZ reorganization, the system locally recovers a scale Λ_3 , roughly equal to the initial spatial scale Λ_1 . (b) In Fourier-space, the initial OD pattern is centered at $k_{\max}\tilde{\mathbf{x}}$. $\delta\Lambda > 0$ equals to a shift $\delta k < 0$. Recovery of the initial spacing corresponds to the growth of new Fourier modes at $(k_{\max} + \delta k)\tilde{\mathbf{x}} \pm q_y\tilde{\mathbf{y}}$ with $((k_{\max} + \delta k)^2 + q_y^2)^{1/2} \approx k_{\max}$. (c) Regions of linear instabilities of the pattern in the $(\delta k, r)$ -plane (redrawn from [29]). The region of the zigzag instability (ZZ, red shaded) is to the left of the $\delta k = 0$ -axis, and is bounded by a parabola. EH (yellow shaded) denotes the regions of the so called Eckhaus instability (see **Methods** and **Supplementary Material**) where a new OD stripe is inserted ($\delta k < -\sqrt{r/3}$) or withdrawn ($\delta k > \sqrt{r/3}$). For $-\sqrt{r} < \delta k < -\sqrt{r/3}$ these two regions overlap (striped region). Note that the ZZ instability occurs for an arbitrary small area increase. Note also that the ZZ instability is the generic behavior for 2-dimensional isotropic dynamics with linear Mexican-Hat type interactions under the influence of isotropic area increase.

Development of OD Columns in the Elastic Network Model Without Growth

The Elastic Network (EN) model [19, 20, 28, 30] describes the development of OD columns by Hebbian plasticity mechanisms with soft competition, as well as lateral intracortical interactions. In this section, we study this model under normal conditions (without growth) and characterize its Turing mechanism.

In the continuous version of the EN model [28], the OD map in V1 at a given time t is described by a real valued field $o(\mathbf{x}, t)$, where $\mathbf{x}$ represents the position on the cortical surface. Negative values of $o(\mathbf{x}, t)$ indicate a preference for input from the ipsilateral eye, positive values a preference for the contralateral eye. This field follows a dynamics

$$\partial_t o(\mathbf{x}, t) = \langle [s_o - o(\mathbf{x}, t)] A_\sigma(\mathbf{x}, \mathbf{S}, o(\cdot, t)) \rangle_{\mathbf{S}} + \eta \Delta o(\mathbf{x}, t), \quad (1)$$

where

$$A_\sigma(\mathbf{x}, \mathbf{S}, o(\cdot, t)) = \frac{e^{-(|\mathbf{s}_r - \mathbf{x}|^2 + |s_o - o(\mathbf{x}, t)|^2)/2\sigma^2}}{\int d^2y e^{-(|\mathbf{s}_r - \mathbf{y}|^2 + |s_o - o(\mathbf{y}, t)|^2)/2\sigma^2}} \quad (2)$$

is the cortical activity pattern, σ controls the receptive field size in the stimulus parameter space, $\langle \cdot \rangle$ denotes the average of the ensemble of visual stimuli $\{\mathbf{S}\}$, η measures the strength of lateral interactions and Δ is the two-dimensional Laplacian. Selectivities $o(\mathbf{x})$ are modified through the cumulative effect of a large number of activity events, evoked by the complete stimulus ensemble. Visual stimuli $\mathbf{S} = (\mathbf{s}_r, s_o)$ are pointlike and characterized by a location $\mathbf{s}_r$ and an OD value s_o , which describes whether the activated units are forced to prefer the ipsilateral ($s_o < 0$) or the contralateral ($s_o > 0$) eye (see **Methods**).

By linear stability analysis, we show that OD columns segregate by a Turing mechanism. We linearize eq. (1) around the homogeneous nonselective state $o(\mathbf{x}) = 0$ (see **Methods**). Due to translation and rotation invariance of the model, the eigenfunctions of this linearized dynamics are Fourier modes $\sim e^{i\mathbf{k}\mathbf{x}}$ with eigenvalues [18, 28]

$$\lambda(k) = -1 + \frac{1}{\sigma^2} \left(1 - e^{-k^2\sigma^2} \right) - \eta k^2 \quad (3)$$

only depending on the absolute value $|\mathbf{k}| = k$ (**Fig. 4a**). The amplitude of any small perturbation containing a Fourier mode with wave vector $\mathbf{k}$ will evolve $\sim \exp(\lambda(k)t)$, and therefore spatial frequencies with $\lambda(k) < 0$ are exponentially damped, whereas those with $\lambda(k) > 0$ grow exponentially (shaded region in **Fig. 4a**). For sufficiently small σ , $\lambda(k)$ exhibits a positive maximum at $r(\sigma, \eta) = \lambda(k_{\max})$, where $k_{\max} > 0$. Thus, the maximum will be at some finite wavelength $\Lambda_{\max} = 2\pi/k_{\max}$ which is characteristic for a Turing instability (see also **Supplementary Material**). We refer to r as the control parameter. Importantly, this defines the timescale $\tau = 1/r$ on which OD columns segregate and the spatial scale $\Lambda_{\max}$ which will be the dominant scale of the emerging pattern of OD columns. Since experiments in cat suggest that the segregation of OD columns takes place within a few days [10, 11], one may identify τ with approximately one day in cat postnatal development. Thus, in the EN model, OD columns segregate because the nonselective state $o(\mathbf{x}) = 0$ becomes unstable and Fourier modes with period $\sim \Lambda_{\max}$ grow exponentially if σ is below a critical value.

Next, we confirmed this by numerical simulations of the full model, eq. (1). To approximate the stimulus ensemble in eq. (1) a large, random sample of pointlike stimuli was drawn at each time step (see **Methods** for details). Different realizations of OD development were obtained by presenting different stimulus samples. In agreement with previous work [19, 20], a structure strongly resembling the OD patterns observed in cat V1 emerges after a few time steps (see **Fig. 1a**, **Fig. 4b**). As a measure of the average cortical selectivity, we monitor the mean field amplitude $\mathcal{A}(t) = \int d^2x |o(\mathbf{x}, t)|$. It shows a rapid increase during the first 10τ and saturates thereafter (**Fig. 4c**). At $t = 4\tau$ the OD pattern already exhibits a visible characteristic spacing which becomes more dominant in the later time course. Confirming our linear stability analysis, we find that the early pattern consists of Fourier modes with wavelength $\sim \Lambda_{\max}$ (**Fig. 4b**, **d**).

We quantify each simulated OD development by the wavelet-method previously applied to our experimental data and compute three parameters characterizing the map layouts. First, we estimate the mean column spacing Λ . We find, that it reaches a constant value after approximately 10τ close to the predicted value $\Lambda_{\max}$ (**Fig. 4d**). Next, we compute the number of hypercolumns N_{HC} in a simulated area. This number changes only very little after OD columns have segregated (**Fig. 4e**). Finally, we calculate the bandedness, α , which quantifies the tendency of OD domains to form elongated parallel stripes (see **Methods** for details). In all our simulations, the initial phase of OD segregation is followed by a long stage of slow rearrangements leading to progressively more stripe-like OD layouts. This behavior is captured by a monotonically increasing bandedness, α , (**Fig. 4f**). In fact, we numerically find that OD stripes, for which the bandedness α is maximal, are stable solutions of the EN model. We do not find any other stable solution.

Zigzag Reorganization in the Elastic Network Model

Next, we directly tested our hypothesis that the EN model shows a ZZ reorganization upon isotropic area increase. As a first step, we considered instantaneous changes in domain sizes. We initiated simulations with a sine wave pattern $\sim \sin(k_{\max}x)$, a state very similar to an attractor of the dynamics (**Fig. 5**). After a brief relaxation period we abruptly changed the size of the simulated system (see **Methods** and **Video 2** in **Supplementary Material**) and studied its subsequent dynamics. **Figure 5** displays snapshots of a typical example of such a simulation ($r = 0.15$, $\eta = 0.025$), with a moderate area increase ($\delta k/k_{\max} = -0.15$). After the size reset at $t = 10\tau$, stripes start to bend sinusoidally (**Fig. 5a**, upper row). In the power spectrum, the growing of additional Fourier modes with wavenumber $\sim (k_{\max} + \delta k)\tilde{\mathbf{x}} \pm q_y\tilde{\mathbf{y}}$ can be observed (**Fig. 5a**, lower row and compare to **Fig. 3b**). These features are typical for




Figure 4: Development of OD column layout in the Elastic Network model without growth. (a) Eigenvalues of the linearized dynamics around the nonselective initial state, $o(\mathbf{x}) = 0$. For $r(\sigma, \eta) > 0$, Fourier modes with positive growth rate form a band around $k_{\max}$, corresponding to an annulus in 2D. OD columns with spatial scale $\Lambda_{\max} = 2\pi/k_{\max}$ segregate on a timescale $\tau = 1/r$ with $r = \lambda(k_{\max})$. (b) Development of OD columns starting from initial condition $o(\mathbf{x}, t = 0) = 0$ ($\eta = 0.025$, $r = 0.2$). At $t = 4\tau$, the emerging pattern already exhibits characteristics of OD columns. After 10τ , columns start to merge and progressively reorganize towards a stripe-like layout. (c-f) Time courses of average OD segregation $\mathcal{A}$ (c) and of three parameters, characterizing OD layouts during development ($N = 50$ realizations, gray: individual traces, black: example from b, red: mean value): (d) mean column spacing Λ (e) number of hypercolumns N_{HC} (f) mean bandedness α . Whereas $\mathcal{A}$, Λ , and N_{HC} reach mature levels around 10τ and exhibit only little variability, the bandedness α increases during the entire time course, expressing the fact that solutions slowly converge towards ideal OD stripes. Furthermore, α -time courses display a large variability, capturing differences between individual realizations of OD map development.

the ZZ instability, thus confirming our hypothesis that the EN model shows ZZ reorganization upon isotropic area expansions. The mean column spacing Λ exhibits a rapid increase at 10τ , induced by the instantaneous area increase, and subsequently decreases almost to the initial value close to $\Lambda_{\max}$ (Fig. 5b). This expresses the fact that the ZZ reorganization restores the column spacing locally. Consequently, the number of hypercolumns N_{HC} in the OD map increases (Fig. 5c). The change in layout of OD columns due to ZZ reorganization is well captured by the drop in the bandedness α (Fig. 5d).

Next, we examined the degree of ZZ reorganization under various instantaneous size increases. We explicitly traced out the regions of stability sketched in Figure 3c for the EN model. We analyzed the time scale τ_{ZZ} on which ZZ reorganization develops, relative to the time scale of column segregation τ , by fitting the ZZ modes in the power spectrum with Gaussians and measuring their peak heights as a function of time. Figure 6a shows a typical example of such a fit ($r = 0.15$, $\delta k/k_{\max} = -0.15$, $t = 110\tau$). The central peak originates from the initial stripe pattern, whereas both of the side peaks are the ZZ modes. To explore the full range of expected rearrangements in our model, we systematically vary the control parameter, r , and the total increase/decrease, δL , of the linear extent, L , equivalent to varying $\delta k/k_{\max}$ (compare to Fig. 3c, see Methods for details). We observe three different types of behavior (Fig. 6c): (i) Even slight area increases result in an observable sinusoidal

rearrangement, which can be identified as the ZZ instability. Small increases in area result in reorganizations on time scales τ_{ZZ} much larger than τ . For considerably large increases, time scales τ_{ZZ} as low as 2τ can be observed. (ii) For very large instantaneous area increases or small control parameters (grey region in Fig. 6c) the initial pattern decays exponentially after size reset and OD segregation restarts with a pattern of characteristic wavelength Λ (pattern plowing). (iii) OD stripes are found to be stable against contraction of the domain ($\delta k > 0$). However, no Eckhaus instability (see Supplementary Material) is observed, even in case of considerable area decrease (Fig. 6c).

In conclusion, this analysis shows that for any arbitrarily small instantaneous area increase ($\delta k < 0$, small), the EN model displays ZZ reorganization, evolving on time scales larger than the time scale τ of OD column segregation. This effect is robust against a variation of r and is, therefore, the generic behavior of the EN model with instantaneous area increase.

Realistic Growth Scenarios

What specific conclusion can be drawn from these theoretical considerations that can be detected experimentally? Evidently, an instantaneous area increase may not provide the best approximation of cortical growth. Moreover, a perfect stripe solution may not be a good approx-




Figure 5: ZZ reorganization in the EN model after instantaneous isotropic area increase. (a) Snapshots of a simulation starting from a near steady state solution of the EN model, a stripe-like OD pattern (upper row) and corresponding power spectra (lower row) ($\eta = 0.025$, $r = 0.15$). At 10τ , the linear extent of the system increased abruptly by 18% ($\delta k/k_{\max} = -0.15$). After instantaneous area increase, OD domains bend sinusoidally and additional modes appear at $(k_{\max} + \delta k)\hat{x} \pm q_y\hat{y}$ in the power spectra. (b-d) Time courses of mean column spacing Λ (b), number of hypercolumns N_{HC} (c), and mean bandedness α (d). At 10τ , Λ increases at first by $\approx 18\%$, but subsequently decreases to almost its initial value. Consequently, N_{HC} increases over the considered period. α shows a characteristic drop, reflecting the bending of OD domains. Note that upon area increase the EN model displays all features typical for ZZ reorganization.




Figure 6: ZZ reorganization in the Elastic Network model under instantaneous isotropic area increase: Timescales and parameter regimes. Compare to the general stability diagram sketched in **Fig. 3c**. (a) Measuring growth rate of zigzag modes. Black crosses mark the power spectrum $P(k_x = k_{\max}, k_y)$. Most power is concentrated around the origin $k_y = 0$, stemming from the initial plane wave, and at two secondary peaks that correspond to the emerging zigzag modes (compare to **Fig. 5**). Red curve is a fit of three Gaussians to the power spectrum (**Methods**). (b) Ratios between the peak height p_c of the central Gaussian and height p_{zz} of the side peaks at different time points (black crosses). Red line marks linear fit to the logarithm of p_{zz}/p_c in the region of exponential growth. The growth rate of zigzag modes is estimated as the slope. In (c), its inverse, the time scale τ_{zz} , is compared to the intrinsic time scale τ of OD segregation for a large set of $(\delta k, r)$ -values. ZZ reorganization was observed for all simulated $r > 0$ and $\delta k < 0$. For small increases in area size, reorganization grows on time scales τ_{zz} much larger than the intrinsic time scale τ . For considerably large area increases, time scales τ_{zz} as low as 2τ are observed. In the grey region, the original pattern decays and a new OD pattern segregates (pattern plowing). Since simulation time diverges for $r \rightarrow 0$ (**Methods**), no simulations were carried out in the white region close to $r = 0$.

imation of the OD layout in cat visual cortex.

Therefore, we next investigated how the ZZ reorganization generalizes to more realistic growth scenarios, a cortex that is growing continuously beginning before the time OD columns segregate. To this end, we performed pairs of simulations, one for which we continuously increased the domain size by various amounts (growing system) and one for which we fixed the domain size after 10τ (non-growing system). To approximate realistic conditions, we initiated our simulations with the uniform and nonselective state $o(\mathbf{x}) = 0$ and linearly increased the linear extent of the simulated regions by 20%, 40%, 60% and 80% between $t = 0\tau$ and $t = 100\tau$ (corresponding to isotropic total area increases of 44%, 96%, 156% and 224%). Test simulations with sublinear and logistic growth did not lead to significantly different results. By measuring average column spacing, Λ , number of hypercolumns, N_{HC} , and bandedness, α , we quantitatively characterized growth-induced changes and reorganizations for complex spatial OD layouts under realistic growth conditions.

Figure 7a shows snapshots of a typical example of these simulation pairs ($r = 0.16$, $\eta = 0.025$, 60% linear extension). Shortly after 10τ , the non-growing system (upper row) tends to converge towards a stripe-like layout. In contrast, the growing system (lower row) exhibits a much more irregular arrangement of columns over the whole time course of the simulation. In fact, the growth induced reorganization of OD layouts shows aspects of the ZZ reorganization. The column spacing of the growing system remains close to the value of the non-growing system (**Fig. 7b**). At the same time, N_{HC} increases dramatically for the growing system (**Fig. 7c**). Comparing the time course of the bandedness α for both simulations (**Fig. 7d**) shows that growth induced a drop in bandedness. Hence, all three parameters show a similar time course as for the stereotypic ZZ reorganization (compare **Fig. 5b-d**). Thus, also for a realistic growth scenario, the growth induced reorganization displays the fundamental characteristics of a ZZ reorganization.

To systematically explore how the reorganization depends on the growth rate and the control parameter, r , we quantify the decrease in bandedness for various r and total size increases (50 simulations per data point) (**Fig. 8**). A polynomial fit to the time courses of the bandedness, α , provides us with two characteristic quantities: the size of the bandedness drop, $\Delta\alpha$, and the duration, Δt , the time interval between the minimum of the drop and the preceding maximum (**Fig. 8a**). Even though the simulations exhibit a large variability, bandedness decrease is consistently observed. **Figure 8b** demonstrates that the percentage of simulations showing a significant bandedness drop (with $\Delta\alpha > 0.05$, $\Delta t > 15\tau$) is largely independent of the control parameter r . With considerable growth, this percentage is close to 100% as compared to $< 6\%$ for non-growing systems. For smaller values of r the drop is more pronounced. Interestingly, the average drop in bandedness depends only weakly on the total

area increase (**Fig. 8c**). Thus, even moderate growth induces substantial reorganization of OD layouts as compared to the non-growing system. Perhaps surprisingly, the duration Δt , i.e. the time for which α monotonically decreases, is largest for small total area increases. (**Fig. 8d**).

Reorganization of OD columns During Growth in Model and Experiment

Finally, we looked for signatures of ZZ reorganization in experiment and compared it to the model predictions. We quantitatively analyzed the 2-DG/proline autoradiographs by assessing column spacing Λ , number of hypercolumns N_{HC} , and bandedness α , and compared the time courses of these values with those obtained in our simulations. As outlined above, our simulations of OD segregation together with experimental data [10, 11] suggest that one model time scale, τ , corresponds to approximately one day in cat postnatal development. In approximate accordance with experimental data from Duffy et al. [3], we used 60% linear extension between $t = 0$ and $t = 100\tau$ corresponding to an area increase of 156%. Similar results were obtained when different growth rates are used (see **Fig. 8**) or when growth was sublinear or logistic (see **Supplementary Material**).

Figure 9a displays time courses of the hypercolumn number N_{HC} in 50 simulations obtained for different stimulus sets (see **Methods** and **Video 3** in **Supplementary Material**). N_{HC} is linearly increasing with simulation time (increase of 47% between $t = 30\tau$ and $t = 70\tau$). **Figure 9d** shows the hypercolumn number N_{HC} for the 2-DG/proline autoradiographs. The number of hypercolumns for kittens at later stages in postnatal development is significantly larger, compared to kittens at early stages ($r = 0.7720$, $p < 0.0002$). At PD28, V1 contains on average 260 ± 40 ($N = 5$) hypercolumns, around PD72 an average of 319 ± 17 ($N = 6$) (increase of 23%) hypercolumns. The increase in N_{HC} in our simulations stems from the fact that the column spacing Λ increases only transiently during continuous isotropic area increase (**Fig. 9b** and **5**). Interestingly, this behavior, characteristic of ZZ reorganization, appears to be present also in the data as a closer look at the 2-DG/proline data set reveals (**Fig. 9e** and **Fig. 2a**). Column spacings are $\approx 5\%$ larger for kittens of intermediate ages between PD40 and PD80 and the time course of column spacing in model and experiment appear very similar. Furthermore, the bandedness α decreases considerably with age, especially between P30 and P60. This decrease in bandedness is highly significant ($r = -0.5754$, $p < 6 \cdot 10^{-5}$) indicating that OD columns while largely preserving their initial spacing reorganize and develop a more isotropic layout over time. As shown in **Figure 8**, such a drop in α is characteristic of a ZZ reorganization. Moreover, as for the column spacing Λ and hypercolumn number N_{HC} , the change in bandedness as well as the time scale on




Figure 7: OD development with growth. (a) Snapshots of OD development in the EN model on a domain growing by 156% between $t = 0$ and $t = 100\tau$ (upper row) and with the growth stopped at $t = 10\tau$ (lower row) ($r = 0.16$, $\eta = 0.025$). (b-d) Time courses of mean column spacing Λ (b), number of hypercolumns N_{HC} (c), and bandedness α (d) for the simulations in a. Despite its large area increase, the growing system displays only a mild and transient increase in Λ , implying a strong increase in N_{HC} . Note that also a continuous increase of area induces reorganization of OD layouts that shows the characteristic features of the zigzag reorganization.

which this changes evolves are in good agreement to the EN model under a realistic growth scenario. We therefore conclude that OD columns reorganize weeks after segregation and that models for their activity-dependent formation suggest that this reorganization is induced by cortical growth.

III. DISCUSSION

In cat visual cortex, cortical columns are present long before its postnatal growth comes to an end. How does cortical growth affect its columnar architecture? During growth, the cell bodies of most neurons are progressively displaced from each other. Since this process inevitably requires the modification of neuronal elements, it might also alter considerably the patterns of cortical activity. Interestingly, the critical period, during which circuits are particularly susceptible to changes of activity patterns, overlaps with the period of strong postnatal growth. It is an appealing hypothesis that cortical circuits remain plastic for an extended period in development in order to facilitate the modification of neuronal circuits to adjust for cortical growth. Analyzing the organization of OD columns in cat primary visual cortex during a period of substantial postnatal growth, we provide evidence for reorganization of columnar layouts long after OD columns have been segregated. Our theoretical analysis shows that the observed reorganization is consistent with a fundamental mechanism of growth induced reorganization that is inherent to a very large class of

models for activity-dependent formation of OD columns. By this mechanism, cortical growth induces a characteristic zigzag-like bending of OD columns which largely restores the spacing of columns. Thus, our empirical findings together with our theoretical analysis suggest that cortical growth accompanied by significant displacement of neuronal cell bodies can induce substantial remodeling of neuronal circuits.

a. Reorganization of OD Columns during Postnatal Growth We visualized the overall layout of OD patterns in V1 of kittens between PD28 and PD98 by 2DG/proline labelling. To minimize genetic variation we included a littermate couple into our dataset. In addition, we performed chronic optical recordings of intrinsic signals to follow OD development in individual animals. We analyzed the layout of OD columns with high precision using a wavelet method [23, 25]. We observed no systematic differences in layout between 2DG and proline labelled columns[25]. In accordance with [3, 12, 14] we found a significant postnatal growth of V1 of about 30% over the considered period (**Fig. 1**), but no corresponding increase in column spacing Λ (**Fig. 2**). This implies an increase in the number of hypercolumns N_{HC} per visual cortex, where a hypercolumn is estimated as a cortical region of size Λ^2 . Since an attachment of columns at the peripheral regions of V1 appears unlikely (see **Supplementary Material**), these findings suggest that the layout of OD columns changes during normal development. In fact, as a signature of ongoing modification of columnar architecture during postnatal development, we found that the bandedness of OD columns, α , displays a




Figure 8: Comprehensive analysis of growth-induced reorganization of OD column layouts in the EN model. (a) Two parameters were used to describe bandedness time course (Fig. 7): Size of bandedness drop $\Delta\alpha$ (difference between the maximal bandedness and the minimal subsequent bandedness) and duration Δt (time interval between these two bandedness extrema). Blue curve is a polynomial fit to the bandedness time course (see **Methods**). Other parameters as in Fig. 7. (b) Percentage of simulations with continuous isotropic area increase at different rates between $t = 0$ and $t = 100\tau$ and different values of r ($N = 50$ simulations for each data point, error bars indicate s.e.m.) that show a significant bandedness drop (for which $\Delta\alpha > 0.05$ and $\Delta t > 15\tau$). For considerable growth, this percentage is close to 100% as compared to $< 6\%$ for non-growing systems. (c-d) Average size $\Delta\alpha$ (c) and duration Δt (d) of significant bandedness drops. For smaller values of r the drop is more pronounced, but no clear dependence on the total area increase was found. Δt is largest for relatively small total area increases. Thus, even moderate growth induces a substantial drop in bandedness α in most simulations.

strong negative correlation with age. Early OD layouts exhibit a stronger tendency to form elongated parallel bands and hence are qualitatively different from mature ones. This decrease in bandedness evolves over a period of weeks, long after OD columns have segregated.

b. A Universal Mode of Reorganization Intrigued by these findings, we studied as a model for the activity-dependent development of cortical circuits the two-dimensional Elastic Network (EN) model. Investigation of a simple toy growth scenario, OD stripes subjected to an instantaneous area increase, revealed that the generic mechanism of growth-induced reorganization in this model is the zigzag reorganization (Fig. 5). OD stripes are found to approximately maintain their intrinsic spacing by bending zigzag-like in response to cortical growth (Fig. 5). This bending occurs on timescales τ_{ZZ} much larger than the time scale τ of OD segregation. Thus, assuming the latter takes a few days, the ZZ reorganization evolves over a few weeks.

Furthermore, we performed a quantitative analysis of OD development in the EN model for realistic growth scenarios. To characterize growth-induced reorganizations, the time course of local column spacing Λ , number of hypercolumns N_{HC} and bandedness α were compared to non-growing systems. We observed three features characteristic for zigzag reorganization: (i) a mild and transient increase in column spacing Λ , (ii) a monotonic increase in hypercolumn number N_{HC} , (iii) a decrease of local anisotropy or bandedness α (Fig. 7, 8). All three features are consistent with our experimental observations (Fig. 9). Indeed, assuming one day in development is approximately equal to τ , the time scale of OD segregation in the model (since OD columns are observed to segregate within a few days [10, 11]), the time course of growth induced reorganization in the model as captured by these parameters exhibits the same characteristic features as the reorganization we observe in experiment. Thus, our theoretical analyses indicate that OD columns may gradually reorganize by a ZZ mechanism in response to cortical growth during postnatal development.

Importantly, we have argued that ZZ reorganization as a mechanism for growth induced reorganization applies not only to the EN model, but generalizes to a large class of models for the formation of OD columns, containing e.g. [15, 16, 17, 18, 19]. The ZZ instability belongs to the class of so-called “long wavelength instabilities” [27, 29], describing reorganizations of patterns on scales much larger than the spacing of the pattern itself (see Fig. 3c). Thus, although the changes in neuronal morphologies across V1 during postnatal growth may be rather subtle on the level of individual neurons, they might cause a remodeling of OD columns which is coordinated over large parts of V1.

c. Relation to Previous Work Several experimental studies in different mammalian species have shown that the cortical surface area as well as the volume increases significantly during postnatal growth [3, 12, 14, 38]. Only very few studies have examined the relation between postnatal area increase and the spacing of cortical columns, and have come to contradictory conclusions. In one study [12], 2-DG labelled OD columns in cat visual cortex between 3 and 10 weeks were analyzed. In line with our results, this study finds no correlation between column spacing Λ and age, despite a considerable size increase of V1. Focusing on an earlier period, a longitudinal optical imaging study of OD columns in single strabismic cat found an increase of OD column spacing between PD27 and PD61 [24] consistent with the slight increase we observe over this period (Fig. 9e). The only theoretical study of OD development during cortical growth [39] focuses on the 1D Swindale model [15]. This study predicts a splitting of OD stripes in response to cortical growth, analogous to the Eckhaus instability discussed in the **Methods** and **Supplementary Material**. However, as acknowledged by the authors themselves, generalizations to two dimensions cannot be derived from this study. Indeed, we show that two-dimensional models for




Figure 9: The reorganization of OD layouts in the EN model under a realistic growth scenario is consistent with the reorganization observed in experiment. **(a-c)** Model: Time courses of hypercolumn number N_{HC} **(a)** column spacing Λ **(b)**, and bandedness α **(c)** ($N = 50$ realizations, isotropic total area increase by 156% between $t = 0$ and $t = 100\tau$, $r = 0.16$, $\eta = 0.025$). Black curve shows the average time course, grey shaded region the SEM. Note the transient increase in column spacing Λ and the typical bandedness drop $\Delta\alpha$ between 30τ and 60τ , both characteristic features of the ZZ reorganization. **(d-f)** Data: hypercolumn number N_{HC} **(d)**, column spacing Λ **(e)**, reproduced from **Fig. 2a**, and bandedness α **(f)** for the same 41 hemispheres. N_{HC} increases significantly between the two groups, centered at PD30 and PD70 ($r = 0.7720$, $p < 0.0002$). Column spacings appear to increase transiently at intermediate ages. The bandedness decreases from an average of 0.15 ± 0.04 ($N = 8$) at PD28 to an average of 0.083 ± 0.02 after PD84 ($N = 6$) and exhibits strong negative correlation with age during this period ($r = -0.5754$, $p < 6 \cdot 10^{-5}$). Thus, assuming $\tau \approx 1$ day (see text), the growth induced reorganization in the model as captured by these parameters evolves on a similar time scale and exhibits the same characteristic features as in experiment.

OD development behave qualitatively differently. In particular, we found no splitting of stripe-like OD domains for isotropic area increases.

d. Remodeling of Ocular Dominance The class of models we propose in this work to explain our empirical observations predicts that OD shifts during normal postnatal development in a sub-population of neurons. Such shifts may require a maximal LGN terminal arbor shift of up to $\Lambda/2 \approx 600 \mu\text{m}$ within one time scale of ZZ reorganization, corresponding to a speed of shift of $\approx 60 \mu\text{m} / \text{day}$. An important question is whether the observed structural changes in *in vivo* plasticity of geniculocortical afferents are sufficient to account for this reorganization. In the normal adult mouse neocortex, structural plasticity due to elongation and retraction of axonal arbors has been observed at a speed of up to $35 \mu\text{m} / \text{day}$ [31]. So far, rapid remodeling of LGN afferents in cat visual cortex in response to short-term monocular occlusion has been demonstrated to occur at velocities up to $2 \text{ mm} / \text{day}$ [32]. This suggests that the morphological changes required by the mode of reorganization described in this paper appear to fall within the range of already observed *in vivo* plastic changes.

An alternative conceivable scenario might be that the observed changes in OD layout during growth is due to a particular pattern of displacement of neurons while their OD is held fixed. This displacement should then be such that the spacing between columns changes only slightly and the bandedness decreases progressively. This could be achieved, for instance, by a displacement in the form of a ZZ-pattern similar to the pattern of reorganization

of OD in the studied model class. However, it is unclear at present what mechanism could give rise to a collective displacement of neurons consistent with our empirical observations.

e. A Further Role for the Critical Period According to our theoretical analyses, OD development in the Elastic Network Model can be divided into three phases: (i) the spontaneous emergence of spatially isotropic OD patterns, occurring on a characteristic time scale τ ($t = 0 - 10\tau$) (ii) growth-induced reorganization ($t = 10 - 100\tau$) and, (iii) a subsequent phase during which the segregated OD patterns slowly reorganize towards stripe-like OD layouts ($t > 100\tau$). This third phase covers a period in cat postnatal development which is not in the range of our data set. To the best of our knowledge, no OD layout similar to the stripe-like stationary states of the EN model has ever been reported in cat V1. Thus, the EN model might not be an appropriate model to describe the layout of cat OD columns during this developmental stage. The first phase is fairly consistent with experimental observations on OD segregation [10, 33] and suggests that τ can be approximately identified with one day in cat postnatal development. Hence, the second phase of growth-induced reorganization in the EN model corresponds to the phase in cat development during which OD columns reorganize. Moreover, this phase largely overlaps with the critical period, as depicted in (**Fig. 1c**).

The impressive ability of cortical circuits to reorganize during the critical period has been demonstrated in numerous studies by artificially manipulating cortical

activity, e.g. by monocular deprivation [34, 35, 36, 37]. Our findings suggest a possible new role of critical period plasticity in normal development. Neuronal cell bodies become progressively displaced from each other during growth, which inevitably requires morphological changes of neuronal elements. Intracortical synaptic connections that are established early in development must be either replaced or the respective dendritic and axonal elements must be elongated. However, this may in turn modify the spatial and temporal properties of dendritic integration and signal transduction between neurons. Since growth induces displacement in a large fraction of pairs of neurons, this could have a severe effect on cortical processing. Critical period plasticity may serve to compensate for these growth induced modifications of neuronal elements and to facilitate the modifications of neuronal circuits necessary to adjust for cortical growth.

IV. METHODS

Optical Imaging Data

Surgery. Anaesthesia was induced with an intramuscular injection of ketamine (10 mg / kg Ketanest®, Parke-Davis, Berlin, Germany) and xylazine hydrochloride (Rompun®, Bayer AG, Leverkusen, Germany) and maintained throughout the experiment using nitrous oxide/oxygen anaesthesia (50% N₂O/50% O₂), supplemented with halothane (0.8-1.2%, Eurim Pharma, Germany). The ECG, pulmonary pressure, end tidal CO₂ (3-4%), and rectal temperature (37-38°) were continuously monitored. The animals head was fixed in a stereotactic frame by means of a metal nut cemented to the skull. For optical imaging of V1 a craniotomy was performed centered at Horsley-Clarke coordinate P4. All experiments were performed when the animals were between 28 and 94 days old. Successive experiments in the same animal were performed with intervals of 7 days and in one case 42 days. All animal experiments have been performed according to the German Law on the Protection of Animals and the corresponding European Communities Council Directive of November 24, 1986 (86/609/EEC).

Visual Stimulation. Animals were stimulated monocularly with high-contrast square-wave gratings (subtending 90°×60° visual field) of four orientations (0°, 45°, 90°, and 135°) moving at a speed of 2cyc/s with a spatial frequency of 0.5cyc/deg. Stimuli were generated by EZV-Stim software (Optical Imaging Inc., Rehovot, Israel) and presented on a LG Electronics Flatron 295 LCD-monitor (luminosity 180 cd/m²; contrast 300:1; refresh rate 85 Hz; resolution 1600×1200 pixel) at a distance of 25 cm. The eyes were treated with atropine and Neosynephrine® and refracted appropriately using corrective corneal contact lenses with artificial pupils with a diameter of 3 mm.

Data Acquisition. The cortical surface was illumi-

nated by means of two adjustable light guides attached to a tungsten-halogen lamp (Spindler & Hoyer, Göttingen, Germany) equipped with interference filters for different wavelengths. The vascular pattern of the cortex was visualized at 546 nm ± 10 nm (green), cortical activity maps at 707 nm ± 1 nm (red). During data acquisition of intrinsic signals, the camera was focused 650–750 μm below the cortical surface. A tandem-lense was used for imaging [44]. The ORA 2001 system (Optical Imaging Inc.), equipped with a cooled Theta CCD system (384×288 pixel chip from Thomson-CSF) was used for collecting the intrinsic signals. We acquired a series of frames every 12 s, whereby a grating of a given orientation was presented for 2s in a static mode, followed by 4.2 s of data acquisition during which the grating was moved in both directions along the axis orthogonal to its orientation. We used episodic stimulation during data acquisition (7 frames of 600 ms duration). The first frames were excluded from further analysis. The stimulus presentation was monocular, an eye shutter was used to conceal the eyes. A single stimulus trial consisted of 2×8 stimulus conditions (4 grating orientations for the left and the right eye) and 8 isoluminant blanks presented in a random sequence. Twelve trials were usually presented to obtain a map, so that every stimulus was shown 24 times. We first calculated “single condition maps” in which the images acquired during presentation of a particular stimulus were divided by the sum of all different stimulus conditions (“cocktail blank procedure”) [45, 46]. Differential maps for ocular dominance (OD) were calculated by summing all left eye activity maps and subtracting all right eye activity maps.

Data Preprocessing. All computed differential maps were preprocessed in order to remove overall variations in signal strength and measurement noise. To achieve this, first the local average signal strength

$$\bar{I}(\mathbf{x}) = \frac{\int_{A17} d^2y K(\mathbf{x} - \mathbf{y}) I_0(\mathbf{y})}{\int_{A17} d^2y K(\mathbf{x} - \mathbf{y})},$$

was calculated using the Fermi filter kernel $K(\mathbf{y}) = 1/(1 + \exp((\mathbf{y}^2 - hp)/\beta))$ and subtracted from the pattern

$$I_1(\mathbf{x}) = I_0(\mathbf{x}) - \bar{I}(\mathbf{x})$$

We used $\beta = 0.2$ and a cutoff wavelengths $hp = 1.8$ mm. Then we lowpass filtered the image with a second Fermi filter using a cutoff wavelengths of $lp = 0.7$ mm and $\beta = 0.2$. The pattern was then centered to yield $\int_{A17} d^2x I(\mathbf{x}) = 0$ and its variance was normalized to one. This overall bandpass filtering ensured that structures on a scale between 0.7 mm and 1.8 mm were only weakly attenuated by the preprocessing and enabled us to do further quantitative analysis.

2 DG - Autoradiography

Animals. We analysed patterns of OD columns in primary visual cortex (V1) of 31 cats (49 hemispheres). Some of the OD maps of these animals have already been published (see [11, 25] and references therein). OD patterns were labelled with either 2- ^{14}C -deoxyglucose (2-DG) autoradiography [47] after monocular stimulation of the animals or by ^3H -proline autoradiography after injection of the labelled proline into one eye which labels the thalamocortical afferents of that eye in cortical layer IV [48, 49]. All experimental procedures are described in detail in [21, 22, 50]. A list of the dataset as well as a description of the preprocessing is given in [25].

Data Analysis Method

Column spacing Λ and bandedness α of both, data and simulations, were analyzed using the wavelet-method introduced in [23].

Estimation of Column Spacing

For each OD pattern $I(\mathbf{x})$ we calculated a wavelet representation, using wavelets which covered only a few hypercolumns but exhibited a strong periodicity. These representations were obtained from

$$\hat{I}(\mathbf{x}, \theta, l) = \int_{A17} d^2y I(\mathbf{y}) \psi_{\mathbf{x}, \theta, l}(\mathbf{y}),$$

where $\mathbf{x}, \theta, l$ are the position, orientation and scale of the wavelet $\psi_{\mathbf{x}, \theta, l}$ and $\hat{I}(\mathbf{x}, \theta, l)$ denotes the array of wavelet coefficients. We used complex Morlet wavelets defined by

$$\psi(\mathbf{x}) = \exp\left(-\frac{(x_1^2 + \sigma^{-2}x_2^2)}{2}\right) e^{i\mathbf{k}_\psi \cdot \mathbf{x}}$$

and

$$\psi_{\mathbf{x}, \theta, l}(\mathbf{y}) = l^{-1} \psi\left(\Omega^{-1}(\theta) \frac{\mathbf{y} - \mathbf{x}}{l}\right),$$

where

$$\Omega(\theta) = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$$

is the two-dimensional rotation matrix.

The characteristic wavelength of a wavelet with scale l is $\Lambda_\psi l$ with $\Lambda_\psi = 2\pi/|\mathbf{k}_\psi|$. σ denotes the anisotropy of the wavelet. We used relatively large ($\mathbf{k}_\psi = (7, 0)$) isotropic wavelets to estimate local column spacing $\Lambda(\mathbf{x})$. We first calculated

$$\bar{I}(\mathbf{x}, l) = \int_0^\pi \frac{d\theta}{\pi} |\hat{I}(\mathbf{x}, \theta, l)|$$

of the wavelet coefficients for every position $\mathbf{x}$ and then determined the scale by computing $\bar{l}(\mathbf{x}) = \text{argmax}_l(\bar{I}(\mathbf{x}, l))$. The corresponding characteristic local wavelength was then $\Lambda(\mathbf{x}) = \bar{l}(\mathbf{x})\Lambda_\psi$. We used 12




Figure 10: Wavelet analysis of columnar layouts. (a) Examples of two wavelets superimposed on a stripe-like region of a simulated OD pattern. The real parts of the complex-valued wavelet are depicted. Positive regions are delineated by red lines, negative regions by green lines. The wavelet of optimal spatial frequency and orientation (solid line) and one of non optimal orientation (dashed line) are shown. (b) Wavelets superimposed on sinusoidally bended stripe patterns, characteristic for the ZZ reorganization. (c-d) The normalized squared modulus of the wavelet coefficients as a function of orientation θ for the pattern in a (c) and for the pattern in b (d). Note that for the stripe-like pattern $|\hat{I}(\mathbf{x}, \theta)|^2$ is strongly modulated and exhibits a pronounced peak. The sinusoidally bended stripes lead to a broader θ -dependence and a less pronounced peak. This difference is captured by the bandedness parameter α .

equally spaced orientations and 16 scales l_j between 0.5 mm and 2.0 mm. $\bar{l}(\mathbf{x})$ was then estimated as the maximum of a polynomial of 6th degree in l , fitting the $\bar{I}(\mathbf{x}, l_j)$ (least square fit) for a given position $\mathbf{x}$. Based on the local column spacing $\Lambda(\mathbf{x})$ we calculated the mean column spacing $\Lambda = \langle \Lambda(\mathbf{x}) \rangle_x$ and the number of hypercolumns in a given area A , $N_{HC} = A/\Lambda^2$. Here and in the following, $\langle \cdot \rangle_x$ denotes the average over V1 or the simulated domain.

Estimation of Local Bandedness α The orientation dependence of the wavelet coefficients was used to calculate a parameter measuring the anisotropy of local pattern elements as shown in Fig. 10. For a pattern consisting of elongated bands, the magnitude of the wavelet coefficients depends strongly on the wavelet orientation and is largest if the orientation of the wavelet matches the orientation of the bands (Fig. 10a, c). For a pattern consisting of more bended stripes or isotropic patches, the wavelet coefficients depend only weakly on the orientation of the wavelet (Fig. 10b, d). Using only wavelets with local wavelength $\Lambda(\mathbf{x})$, we therefore calculated

$$s'(\mathbf{x}) = \int_0^\pi d\theta |\hat{I}(\mathbf{x}, \theta)|^2 e^{i2\theta} / \int_0^\pi d\theta |\hat{I}(\mathbf{x}, \theta)|^2. \quad (4)$$

We defined the local bandedness as

$$s(\mathbf{x}) = \int_{V_1} d^2y K(\mathbf{x}-\mathbf{y})s'(\mathbf{y}) \Big/ \int_{V_1} d^2y K(\mathbf{x}-\mathbf{y}) , \quad (5)$$

where $K(\mathbf{x}) = \frac{1}{2\pi\sigma^2} \exp(-\mathbf{x}^2/2\sigma^2)$ and $\sigma = 1.3\Lambda$. To estimate (4) and (5), wavelet coefficients were computed for 9 equally spaced orientations. With this choice of σ , $s(\mathbf{x})$ is sensitive to the occurrence of band-like regions that extend over the size of a hypercolumn. Based on the local bandedness $s(\mathbf{x})$ we computed the bandedness $\alpha = \langle s(\mathbf{x}) \rangle_x$ to characterize the overall layout of simulated and measured OD patterns.

Parameter Range of ZZ instability

The stability diagram shown in **Fig. 3c** is very general, applying to the class of two dimensional translation and rotation symmetric systems in which the pattern forming process is based on a Turing-mechanism [27]. Assuming a plane wave solution $A(x, y, t)e^{ik_{\max}x}$, close to the Turing-instability its behavior is governed by the Newell-Whitehead equation [51]

$$\partial_t A = rA + (\partial_x - i\partial_{y^2})^2 A - |A|^2 A, \quad (6)$$

where r is called the control parameter of the system. The form of eq. (6) is independent of the microscopic details of the instability mechanism [27]. Inserting the ansatz $A = |A|e^{i(\delta kx + \phi_0)}$, yields a uniform solution $|A| = \sqrt{r - \delta k^2}$ as long as $|\delta k| < \sqrt{r}$. The wavenumber of this solution differs by an amount δk from $k_{\max}$, i.e., their characteristic spacing is either smaller or larger than Λ . Linear stability analysis yields the behavior of this solution depending on the control parameter r and the difference in wave vector δk [52]. Considering general perturbations of the form $\delta A \sim e^{iq_y y} e^{iq_x x}$, the uniform solution is found to be unstable against perturbations with $q_x = 0$ as soon as

$$\delta k < 0.$$

This defines the domain of the ZZ instability (red shaded regions **Fig. 3c**). In contrast, for $q_y = 0$ the uniform solution is only unstable if $|\delta k| > \sqrt{r/3}$. This is the domain of the so-called Eckhaus instability (EH; yellow shaded regions in **Fig. 3c**; see **Supplementary Material** for an illustration). For $-\sqrt{r} < \delta k < -\sqrt{r/3}$ EH and ZZ overlap (striped region in **Fig. 3c**) and model specific properties determine which of the two behaviors is observed in this regime. For $|\delta k| > \sqrt{r}$, eq. (6) has no uniform stationary solution. In this regime, the initial stripe pattern decays exponentially and OD segregation restarts (*pattern plowing*).

Linearization of the Elastic Network Model

First, we demonstrate that the homogeneous nonselective state $o_{\text{hom}}(\mathbf{x}) \equiv 0$ is a stationary solution of eq. (1)

[28]. Inserting $o_{\text{hom}}(\mathbf{x}) \equiv 0$ into eq. (2) yields

$$A_\sigma(\mathbf{x}, \mathbf{S}, o_{\text{hom}}) = \frac{e^{-\frac{|\mathbf{s}_r - \mathbf{x}|^2}{2\sigma^2}}}{2\pi\sigma^2}.$$

After averaging over the ensemble of stimuli with normalized uniform densities ρ_r and ρ_{s_o} , we obtain

$$\partial_t o(\mathbf{x})|_{o_{\text{hom}}} = \int ds_r \rho_r \frac{e^{-\frac{|\mathbf{s}_r - \mathbf{x}|^2}{2\sigma^2}}}{2\pi\sigma^2} \int ds_o \rho_{s_o} s_o = 0.$$

This shows that $o_{\text{hom}}(\mathbf{x}) \equiv 0$ is a fixed point of the dynamics in eq. (1). The stability of this fixed point can be studied by linearizing the r.h.s of eq. (1) [18, 28],

$$\mathcal{F}[o, \mathbf{x}, \sigma, \mathbf{S}] \approx \mathcal{F}[o_{\text{hom}}, \mathbf{x}, \sigma, \mathbf{S}] + \int d^2y \frac{\delta \mathcal{F}}{\delta o(\mathbf{y})} \Big|_{o_{\text{hom}}} o(\mathbf{y}).$$

Assuming a uniform stimulus density ρ_{s_r} across the cortical surface this results in

$$\partial_t o(\mathbf{x}) = \left(\eta \Delta + \frac{\langle s_o^2 \rangle}{\sigma^2} - 1 \right) o(\mathbf{x}) - \frac{\langle s_o^2 \rangle}{4\pi\sigma^4} \int d^2y e^{-\frac{(\mathbf{x}-\mathbf{y})^2}{4\sigma^2}} o(\mathbf{y}). \quad (7)$$

Note that the linearization is only governed by the *variance* of the stimulus ensemble, and not by higher order statistical moments and is invariant under spatial rotations and translations. Due to the latter, its eigenfunctions are Fourier modes $\sim e^{i\mathbf{k}\mathbf{x}}$ and their eigenvalues $\lambda(k)$ only depend on the absolute value $k = |\mathbf{k}|$. By Fourier-transforming eq. (7) with $\langle s_o^2 \rangle = 1$ we obtain $\lambda(k)$ in eq. 3.

The homogeneous nonselective state $o_{\text{hom}}(\mathbf{x}) \equiv 0$ is unstable, if some eigenvalue with $k > 0$ is larger than zero. For $\eta > 0$ and $\sigma > 0$, $\lambda(k)$ has a single maximum at $k_{\max} = \frac{1}{\sigma} \sqrt{\ln(1/\eta)}$. $\lambda(k_{\max})$ is *positive* if the width σ of the activation function is smaller than $\sigma^* = \sqrt{1 - \eta + \eta \ln \eta}$. In this regime, the homogeneous nonselective state is unstable and the maximum positive eigenvalue $r = \lambda(k_{\max})$ defines a timescale $\tau = 1/r$ on which OD columns segregate and a spatial scale $\Lambda_{\max} = 2\pi/k_{\max}$ that approximately corresponds to the spacing of columns.

Numerical Integration

Simulations were performed on a 64×64 grid with periodic boundary conditions. Simulated systems were spatially discretized with at least 4 grid points per $\Lambda_{\max}$ to achieve sufficient resolution in space. Test simulations with larger grid sizes (128×128, 256×256) did not lead to significantly different results. Progression of time was measured in units of the intrinsic timescale τ . Here, the integration time step δt is bounded by the relevant decay time constant of the Laplacian in eq. (1) around $k_{\max}$ and by the intrinsic timescale τ of the system. We used

$\delta t = \min \{1/(20\eta k_{\max}^2), \tau/10\}$ to ensure good approximation to the temporally continuous changes of the OD patterns. Note that, since $\tau = 1/r$, simulation time diverges for $r \rightarrow 0$. The EN model dynamics was simulated using an Adams-Bashforth scheme for the first term on the r.h.s. of eq. (1). The second term was treated by spectral integration, exhibiting unconditional numerical stability. The stimulus parameters s_r and s_o were chosen to be uniformly distributed with densities ρ_r and ρ_{s_o} such that $\langle s_o^2 \rangle = 1$. The stimulus average in eq. (1) was approximated by choosing a random representative sample of N_s stimuli at each integration time step, with

$$N_s = \frac{N_0 \Gamma^2}{\delta t} \varepsilon_s^{-n} \sqrt{\tau g_{00}},$$

where n is the dimensions of the feature space (in our case, $n = 3$), $\Gamma^2 = (L/\Lambda_{\max})^2$ the size of the simulated system in units of $\Lambda_{\max}^2$, ε_s the resolution in feature space, N_0 the number of stimuli we required to sufficiently approximate the cumulative effect of the ensemble of stimuli within each feature space voxel, and $g_{00} = \frac{9(\eta-1)^4}{40\sigma^6}$ a factor that depends on the specific form of nonlinear competition between Fourier modes in the EN model (unpublished data and personal communication from M. Huang and F. Wolf). $\sqrt{\tau g_{00}}$ is proportional to the inverse of the expected mean amplitude of the OD pattern (unpublished data). With $N_0 = 50$ and $\varepsilon_s = 0.15$ we ensured a low amplitude to noise ratio for all the simulations. Typical values for N_s were between 40000 and 200000. To model development prior to OD segregation, we initialized simulations with $o(\mathbf{x}, t = 0) = 0$. Different realizations of OD development were obtained by using different stimulus samples.

Instantaneous Area Increases. We started simulations with a predefined system length of L_0 and used a parameter dependent sine-wave initial condition $1/\sqrt{\tau g_{00}} \sin(k_{\max} x)$ with periodicity $\Lambda_{\max} = 2\pi/k_{\max}$ and amplitude close the amplitude of the expected stationary state. The system was integrated keeping the length of the system fixed for the first 10τ . Within this period the amplitude $\mathcal{A}(t) = \int d^2x |o(\mathbf{x}, t)|$ of the OD stripe patterns relaxed towards a stationary value while OD layout underwent only minor random fluctuations. Next we rescaled the system length to $L_0 + \delta L$ (no change in number of grid points), where

$$\delta L = \frac{(\delta k/k_{\max})}{(\delta k/k_{\max}) - 1} L_0$$

is determined from the desired value of $\delta k/k_{\max}$, thereby increasing ($\delta L > 0$) or decreasing ($\delta L < 0$) the total area

size covered by the OD pattern. Accordingly, we adjusted the number of stimuli, N_s , and, since $\Delta \sim 1/L^2$, the matrix for the spectral integration step. Note that the numerical value of σ , i.e. the width of the activation function $A_\sigma(\mathbf{x}, \mathbf{S}, o(\cdot, t))$, remains unchanged by this rescaling, but its extension relative to L decreases (and the number of grid points per activation blob decreases as well). After rescaling, we continued simulations for another 190τ .

To estimate time scales τ_{ZZ} of the ZZ reorganization, we monitored the growth of the two ZZ modes over time (see **Fig. 6a, b**). At each time point, we fitted Gaussians of variable size, width and position to the original mode and the two growing ZZ modes along the expected axis of the ZZ modes in Fourier space (least square fit) (see **Fig. 6a**). The timescale of ZZ reorganization was extracted by linear fitting (least square fit) the logarithm of the ratio between the peak height of the ZZ modes, p_{ZZ} , and the central peak height stemming from the original pattern, p_c (see **Fig. 6b**). We constraint the fit to values between 15% and 60% of the maximal peak ratio to capture best the regime of exponential growth of the ZZ modes. The fitting procedure was only applied to simulations with maximal peak ratio of at least 0.05.

Continuous Area Increases. To approximate realistic growth conditions, we linearly increased the linear extent L of the simulated regions between $t = 0\tau$ and $t = 100\tau$ (see also **Supplementary Material**). System length L , the Laplacian Δ , and the number of stimuli N_s were updated at each integration step. After 100τ , integration was continued for another 100τ with fixed linear extent. To quantify the size of the bandedness drop $\Delta\alpha$ due to growth as well as the time Δt over which α decreased, we fitted an 8th-order polynomial to each α -time series (least square fit). This captured the coarse grained development but smoothed out fluctuations on intervals smaller than 3τ (see **Fig. 8a**). We only considered bandedness decreases of minimum size $\Delta\alpha > 0.05$ and duration $\Delta t > 15\tau$ in our analysis.

Statistics. r-values denote Pearson's linear correlation coefficient; p-Values were obtained with Student's t-tests. All error bars and $\pm$ are s.e.m..

Acknowledgements

We express our gratitude to F. Wolf for inspiring discussions. We thank M. Huang and F. Wolf for comments on numerical procedures. We thank P. Mehta and S. Palmer for helpful comments on the manuscript.

[1] Dekaban, A.S. & Sadowksy, D. Changes in brain weight during the span of a human life: relation of brain weight to body heights and body weights. *Ann. Neurol.* **4**, 345–

356 (1978).
 [2] Villabanca, J.R., Schmanke, T.D., Crutcher, H.A., Sung, A.C., & Tavabi, K. The growth of the feline brain from

- fetal into adult life i. a morphometric study of neocortex and white matter. *Dev. Brain Res.* **122**, 11–20 (2000).
- [3] Duffy, K.R., Murphy, K.M., & Jones, D.G. Analysis of the postnatal growth of visual cortex. *Vis. Neurosci.* **15**, 831–839 (1998).
 - [4] Cragg, B.G. The development of synapses in the visual system of the cat. *J. Comp. Neurol.* **160**, 147–166 (1975).
 - [5] Purves, D. Neural activity and the growth of the brain. (ed. Brozolo, L.) (Cambridge University Press, 1994).
 - [6] London, M. & Häusser, M. Dendritic computation. *Ann. Rev. Neurosci.* **28**, 503–32 (2005).
 - [7] Martin, K.A. & Whitteridge, D. Form, function and intracortical projections of spiny neurones in the striate visual cortex of the cat. *J. Physiol.* **353**, 463–504 (1984).
 - [8] Häusser, M., Spruston, N. & Stuart, G.J. Diversity and dynamics of dendritic signaling. *Science* **290**, 739–44 (2000).
 - [9] Dan, Y. & Poo, M.M. Spike timing-dependent plasticity of neural circuits. *Neuron* **44**, 23–30 (2004).
 - [10] Crair, M.C., Horton, J.C., Antonini, A., & Stryker, M.P. Emergence of ocular dominance columns in cat visual cortex by 2 weeks of age. *J. Comp. Neurol.* **430**, 235–249 (2001).
 - [11] Rathjen, S. & Löwel, S. Early postnatal development of functional ocular dominance columns in cat primary visual cortex. *Neuroreport* **11**, 2363–7 (2000).
 - [12] Rathjen, S. Schmidt, K.E., & Löwel, S. Postnatal growth and column spacing in cat primary visual cortex. *Exp. Brain Res.* **149**, 151–158 (2003).
 - [13] Sireteanu, R. & Maurer, D. The development of the kitten's visual field. *Vis. Res.* **22**, 1105–11 (1982).
 - [14] Rathjen, S. *et al.* The growth of cat cerebral cortex in postnatal life: a magnetic resonance imaging study. *Eur. J. Neurosci.* **18**, 1797–1806 (2003).
 - [15] Swindale, N.V. A model for the formation of ocular dominance stripes. *Proc. Royal Soc. London, B* **208**, 243–264 (1980).
 - [16] Miller, K.D., Keller, J.B. & Stryker, M.P. Ocular dominance column development: Analysis and simulation. *Science* **245**, 605–615 (1989).
 - [17] Obermayer, K., Blasdel, G., & Schulten, K. Statistical-mechanical analysis of self-organization and pattern formation during the development of visual maps. *Phys. Rev. A* **45**, 7568–7589 (1992).
 - [18] Scherf, O., Pawelzik, K., Wolf, F., & Geisel, T. Theory of ocular dominance pattern formation. *Phys. Rev. E* **59**, 6977–6993 (1999).
 - [19] Goodhill, G.J. & Cimoneriu, A. Analysis of the elastic net model applied to the formation of ocular dominance and orientation columns. *Network (Bristol, England)* **11**, 153–168 (2000).
 - [20] Durbin, R. & Mitchison, G. A dimension reduction framework for understanding cortical maps. *Nature* **343**, 644–647 (1990).
 - [21] Löwel, S. & Singer, W. Monocularly induced 2-deoxyglucose patterns in the visual cortex and lateral geniculate nucleus of the cat. II. awake animals & strabismic animals. *Eur. J. Neurosci.* **5**, 857–869 (1993).
 - [22] Löwel, S. & Singer, W. Monocularly induced 2-deoxyglucose patterns in the visual cortex and lateral geniculate nucleus of the cat. I. anaesthetized and paralysed animals. *Eur. J. Neurosci.* **5**, 846–856 (1993).
 - [23] Kaschube, M., Wolf, F., Geisel, T., & Löwel, S. Genetic influence on quantitative features of neocortical architecture. *J. Neurosci.* **22**, 7206–7217 (2002).
 - [24] Müller, T. *et al.* An analysis of orientation and ocular dominance patterns in the visual cortex of cats and ferrets. *Neur. Comp.* **12**, 2573–2595 (2000).
 - [25] Kaschube, M. *et al.* The pattern of ocular dominance columns in cat primary visual cortex: intra- and interindividual variability of column spacing and its dependence on genetic background. *Eur. J. Neurosci.* **18**, 3251–3266 (2003).
 - [26] Turing, A. The chemical basis of morphogenesis. *Phil. Trans. Royal Soc. London, B* **237**, 37–72 (1952).
 - [27] Cross, M.C. & Hohenberg, P.C. Pattern formation outside of equilibrium. *Rev. Mod. Phys.* **65**, 854–1086 (1993).
 - [28] Wolf, F. & Geisel, T. Spontaneous pinwheel annihilation during visual development. *Nature* **395**, 73–78 (1998).
 - [29] Manneville, P. *Dissipative Structures And Weak Turbulence*. (ed. Araki H., Libchaber A. & Parisi, G.) (Academic Press, 1990).
 - [30] Erwin, E., Obermayer, K., & Schulten, K. Models of orientation and ocular dominance columns in the visual cortex: a critical comparison. *Neur. Comp.* **7**, 425–468 (1995).
 - [31] De Paola, V. *et al.* Cell type-specific structural plasticity of axonal branches and boutons in the adult neocortex. *Neuron* **49**, 861–875 (2006).
 - [32] Antonini, A. & Stryker, M. P. Rapid remodeling of axonal arbors in the visual cortex. *Science* **260**, 1819–1821 (1993).
 - [33] Crowley, J. & Katz, L.C. Early development of ocular dominance columns. *Science* (2000).
 - [34] Hubel, D.H. & Wiesel, T.N. The period of susceptibility to the physiological effects of unilateral eye closure in kittens. *J. Physiol.* **206**, 419–436 (1970).
 - [35] Shatz, C.J. & Stryker, M.P. Ocular dominance in layer iv of the cat's visual cortex and the effects of monocular deprivation. *J. Physiol.* **281**, 267–83 (1978).
 - [36] Antonini, A. & Stryker, M.P. Plasticity of geniculocortical afferents following brief or prolonged monocular occlusion in the cat. *J. Comp. Neurol.* **369**, 64–82 (1996).
 - [37] Hensch, T.K. Critical period plasticity in local cortical circuits. *Nat. Rev. Neurosci.* **6**, 877–888 (2005).
 - [38] Drenhaus, U., Rager, G., Egli, P. & Kretz, R. On the postnatal development of the striate cortex (v1) in the tree shrew (*tupaia belangeri*). *Eur. J. Neurosci.* **24**, 479–490 (2006).
 - [39] Oster, A.M. & Bressloff, P.C. A developmental model of ocular dominance column formation on a growing cortex. *Bull. Math. Biol.* **68**, 73–98 (2006).
 - [40] Marotte, L.R. Goldfish retinotectal system: continuing development and synaptogenesis. *J. Comp. Neurol.* **193**, 319–334 (1980).
 - [41] Raymond, P.A. & Easter, S.S. Postembryonic growth of the optic tectum in goldfish. i. location of germinal cells and numbers of neurons produced. *J. Neurosci.* **3**, 1077–1091 (1983).
 - [42] Easter, S.S. & Stuermer, C.A. An evaluation of the hypothesis of shifting terminals in goldfish optic tectum. *J. Neurosci.* **4**, 1052–1063 (1984).
 - [43] Constantine-Paton, M. & Law, M.I. Eye-specific termination bands in tecta of three-eyed frogs. *Science* **202**, 639–41 (1978).
 - [44] Ratzlaff, E.H. & Grinvald, A. A tandem-lens epifluorescence microscope: Hundred-fold brightness advantage for wide field imaging. *J. Neurosci.* **36**, 127–137 (1991).

- [45] Bonhoeffer, T. & Grinvald, A. Optical imaging based on intrinsic signals: The methodology. in *Brain Mapping. The Methods* (ed. Toga, A., Mazziotta, J.) (Academic Press, 1996).
- [46] Engelmann, R., Crook, J.M. & Löwel, S. Optical imaging of orientation and ocular dominance maps in area 17 of cats with convergent strabismus. *Vis. Neurosci.* **19**, 39–49 (2002).
- [47] Sokoloff, L. *et al.* The [14c]deoxyglucose method for the measurement of local cerebral glucose utilization: theory, procedure, and normal values in the conscious and anesthetized albino rat. *J. Neurochem.* **28**, 897–916 (1997).
- [48] Grafstein, B. Transneuronal transfer of radioactivity in the central nervous system. *Science* **172**, 177–179 (1971).
- [49] Wiesel, T.N., Hubel, D.H., & Lam, D.M. Autoradiographic demonstration of ocular-dominance columns in the monkey striate cortex by means of transneuronal transport. *Brain Res.* **79**, 273–279 (1974).
- [50] Löwel, S. Ocular dominance column development: Strabismus changes the spacing of adjacent columns in cat visual cortex. *J. Neurosci.* **14**, 7451–7468 (1994).
- [51] Newell, A.C. & Whitehead, J.A. Finite bandwidth, finite amplitude convection. *J. Fluid Mech.* **38**, 279–303 (1969).
- [52] Busse, F. & Whitehead, J.A. Instabilities of convection rolls in a high prandtl number fluid. *J. Fluid Mech.* **47**, 305–320 (1971).
- [53] LeVay, S., Stryker, M.P. & Shatz, C.J. Ocular dominance columns and their development in layer iv of the cat's visual cortex: a quantitative study. *J. Comp. Neurol.* **179**(1), 223–44 (1978).
- [54] Luskin, M.B. & Shatz, C. Studies of the earliest generated cells of the cat's visual cortex: Cogeneration of subplate and marginal zones. *J. Neurosci.* **5**(4), 1062–1075 (1985).
- [55] Luskin, M.B. & Shatz, C.J. Neurogenesis of the cat's primary visual cortex. *J. Comp. Neurol.* **242**(4), 611–31 (1985).
- [56] Crampin, E.J., Gaffney, E.A., & Maini, P.K. Reaction and diffusion on growing domains: Scenarios for robust pattern formation. *Bull. Math. Biol.* **61**(6), 1093–1120 (1999).

V. SUPPLEMENTARY MATERIAL

Adding OD columns at the periphery of V1

At a first glance, a different scenario of postnatal reorganization during V1 growth seems conceivable which assumes the addition of new functional modules at the periphery, thereby preserving the early OD layout in the central regions. Here we illustrate that this scenario in fact implies substantial large-scale rearrangements of OD columns and is therefore rather implausible.

During the period of postnatal growth considered in this paper, the number of neurons remains largely constant [4]. The increase in total area is largely due to the generation of glial cells, the addition of more vasculature and connective tissue, and the myelination of axons, but also involves an increase in neuropil, i.e. the outgrowth and elaboration of axonal and dendritic processes and extensive remodeling of synaptic connections [4, 5, 53, 54, 55]. **Figure 11** schematically demonstrates that under these assumptions, the addition of new OD columns at peripheral regions of V1 while preserving the OD layout in central regions of V1 implies substantial remodelling of OD in large parts of the neuronal population.

Ocular dominance columns can already be visualized at postnatal week 2 in cat V1 [10]. This layout of alternating domains is sketched in the upper row of **Figure 11**. The lower row displays the potential layout of OD domains in the adult animal. Additional columns are inserted at peripheral regions. The column spacing Λ is the same for both OD patterns. Furthermore, the central part of the early and mature OD layout are identical (dashed lines). However, neurons labelled with a red number in the mature animal have changed their preferred eye. Note their wide distribution across the cortical surface. Note further that during the continuous expansion of V1, eye preference changed multiple times in individual termination zones, in particular at the peripheral regions (e.g. regions 1,18). Thus, we conclude that this mode of postnatal reorganization is implausible to occur during normal development.

Eckhaus Instability

The Eckhaus instability is a mode of reorganization predicted to occur for models with effective Mexican-Hat interactions in isotropically growing as well as shrinking systems. Rearrangements of domains consist either in inserting new domains (area increase) or removing existing ones (area decrease). Even if it takes place in a two-dimensional system, it is effectively one-dimensional reorganization. It does not require rotation symmetry of the Turing Mechanism.

The result of an Eckhaus instability can be best understood by considering a simple pattern of OD columns consisting of alternating stripes (**Figure 12**). After

an isotropic area increase, the spacing between the OD stripes is larger than the spacing set by the Mexican-Hat. Subsequently, new stripes of OD domains emerge between existing ones by which the original spacing is eventually recovered (**Fig. 12a**). **Figure 12b** illustrates this in Fourier space. The initial stripe pattern is described by a wave $\sim e^{ik_{\max}\tilde{\mathbf{x}}}$ with spatial wavenumber $k_{\max} = 2\pi/\Lambda_{\max}$. Upon area increase, its wave vector differs by an amount $\delta k < 0$ from $k_{\max}$. Two Fourier modes grow with wave vector $\pm k_{\max}\tilde{\mathbf{x}}$, representing the new OD pattern in **Figure 12b** with similar spacing as the original pattern.

Similarly, after an isotropic area decrease, the spacing between the OD stripes is smaller than the spacing set by the Mexican-Hat. Subsequently, a fraction of the OD stripes is removed and the original spacing is eventually recovered (**Fig. 12c**). **Figure 12d** illustrates this in Fourier space. Upon area decrease, the OD pattern's wave vector differs by an amount $\delta k < 0$ from $k_{\max}$. Two Fourier modes grow with wave vector $\pm k_{\max}\tilde{\mathbf{x}}$, representing the new OD pattern in **Figure 12c** with similar spacing as the original pattern. As shown in our manuscript the domain of the Eckhaus instability in the $(r, \delta k)$ -plane is $|\delta k| > \sqrt{r/3}$ (see **Figure 3c** in the manuscript). For $\delta k < 0$, this domain largely overlaps with the domain of the ZZ instability. Since we did not observe signatures of the Eckhaus reorganization in any of our simulations, we conclude that this mode of reorganization is not a generic mechanism in the Elastic Network (EN) model.

ZZ reorganization of OD stripes during continuous area increase

In our manuscript, we analyzed growth-induced ZZ reorganization of OD stripes in the EN model for instantaneous area increases. Here we point out that the characteristic sinusoidal bending of OD domains during this universal mode of reorganization is also observed for stripe-like OD patterns subject to continuous area increase (**Fig. 13**). We initialized our simulations with a sinusoidal pattern $o(\mathbf{x}) \sim \sin(k_{\max}x)$. After a brief relaxation period of 10τ we linearly increased the linear extent of the simulated regions by 20%, 40%, 60% and 80% between $t = 10\tau$ and $t = 110\tau$ (corresponding to isotropic total area increases of 44%, 96%, 156% and 224%). Columnar layouts generally showed reorganization on similar timescales and the reorganization was more irregular when compared to the scenario of instantaneous area increases (**Fig. 13c, d**). In all simulations, column spacing Λ exhibited a transient increase before the onset of reorganization followed by a decreased towards the initial value (data not shown). Bandedness α decreased after onset of reorganization (data not shown) as expected from the characteristic sinusoidal bending of OD domains (**Fig. 13**). Note that stripe-like OD patterns subject to continuous area increase provide an intermediate scenario between OD stripes subject to in-




Figure 11: Adding columns at the periphery requires large-scale remodelling. Schematic cross-sections through the cortical surface at an early postnatal stage (upper row) and an adult stage (lower row) are sketched. Dark/white circles denote regions with ipsilateral/contralateral eye preference. Note that the number of neurons and the spacing of OD columns is assumed to be constant. In regions labelled with a red number, OD is required to change in order to preserve the initial OD layout in the central part. Note that this scenario implies large-scale rearrangements, in particular at peripheral regions of V1, and for some neurons multiple changes in OD over time.




Figure 12: Eckhaus reorganization upon large isotropic area increase (a, b) or area decrease (c, d). Upon area increase/decrease, OD stripes recover their initial spacing Λ_1 by insertion/removal of OD stripes. (a) During isotropic area increase, the OD pattern transiently acquires a spacing $\Lambda_2 = \Lambda_1 + \delta\Lambda$, with $\delta\Lambda > 0$. By insertion of a new stripe the system recovers a spacing Λ_3 , roughly equal to Λ_1 . (b) In Fourier-space, the initial pattern is centered at $\pm k_{\max}\hat{x}$. $\delta\Lambda > 0$ corresponds to a shift of $\delta k < 0$. The initial spacing is recovered by the growth of new Fourier modes close to modes of the original pattern $\pm k_{\max}\hat{x}$. (c) During isotropic area decrease, the OD pattern transiently acquires a spacing $\Lambda_2 = \Lambda_1 + \delta\Lambda$, with $\delta\Lambda < 0$. By removal of stripe the system recovers a spacing Λ_3 , roughly equal to Λ_1 . (d) In Fourier-space, $\delta\Lambda < 0$ corresponds to a shift of $\delta k > 0$, with $\delta\Lambda > 0$. The initial spacing is recovered by the growth of new Fourier modes at the center of the original pattern, i.e. $\pm k_{\max}\hat{x}$. Note that insertion/removal of columns requires strong area increase/decrease, a regime not expected to be relevant during continuous growth.

stantaneous area increase and realistic growth scenarios for complex OD patterns similar to the OD layouts in young kittens.

Different Scenarios for Realistic Cortical Growth

We tested three different simulation protocols to approximate the postnatal growth of cat V1 (see Fig. 14a, b): (i) linear increase of the linear extent with slope α , starting from L_0 ,

$$L(t) = L_0 + \alpha t,$$




Figure 13: ZZ reorganization of a stripe-like OD pattern in EN model during continuous isotropic area increase. **(a, b)** ZZ reorganization far from Turing Instability ($r = 0.18$, $\eta = 0.025$, total area increase 196% between $t = 10\tau$ and $t = 110\tau$). Until $t = 50\tau$ the OD stripes remain almost unchanged. Subsequently, ZZ reorganization is apparent from the sinusoidal bending of domains **(a)** and the growth of Fourier Modes at $k_{\max}\hat{x} \pm q\hat{y}$ **(b)** **(c)** ZZ reorganization close to Turing Instability ($r = 0.04$, other parameters as in **a**). Soon after the onset of ZZ reorganization the pattern acquires an irregular layout. **(d)** Power spectra for the OD layouts in **c**. Note, that despite rather complex reorganization, weak signatures of ZZ modes can still be identified.

(ii) sublinear increase of the linear extent L , such that the simulated area A increases linearly with slope β ,

$$L(t) = \sqrt{L_0^2 + \beta t},$$

(iii) and logistic increase [56] of the linear extent L

$$L(t) = L_0 \frac{e^{\varepsilon t}}{1 + \xi^{-1}(e^{\varepsilon t} - 1)} \quad \lim_{t \rightarrow \infty} L(t) = \xi L_0.$$

In every protocol, we increased the linear extent of the simulated region by a fixed amount (20%, 40%, 60% and 80%) between $t = 0\tau$ and $t = 100\tau$. This corresponds to isotropic total area increases of 44%, 96%, 156% and 224%. Note that according to [3] the area of cat V1 increases between week 1 and week 12 by approximately 150% percent. Simulations of OD development with different growth protocols yielded qualitatively similar results. Linear and sublinear increases described virtually identical growth scenarios (**Fig. 14a, b**) and provided a reasonable fit to the data on V1 growth shown in [3]. Our systematic quantitative characterization of OD development in the EN model with growth (**Figure 7, 8, 9**

in the manuscript) was carried out based on the scenario of linear increase of the linear extent (red traces in **Fig. 14b**).




Figure 14: Three different simulation protocols of realistic cortical growth. **(a-b)** Time courses of the linear extent L/L_0 **(a)** and the area size A/A_0 **(b)** for linear increase of area (green traces), linear (red traces) increase of the linear extent and logistic growth (blue traces). Simulations of OD development for all three growth protocols yielded qualitatively similar time courses of column spacing Λ , number of hypercolumns N_{HC} and bandedness α . Linear area increase and linear increase of the linear extent are virtually identical and both provide a reasonable fit to the data on cat V1 growth in [3]. Our quantitative characterization of OD development in the EN model with growth (**Figure 7, 8, 9**) was carried out using linear increase of the linear extent (red traces).

Videos

Video 1. Development of OD column layout in the EN model without growth, starting from initial condition $o(\mathbf{x}, t = 0) = 0$ ($\eta = 0.025$, $r = 0.16$). The video displays OD maps (upper left) as well as time courses of the parameters mean column spacing Λ (upper right), number of hypercolumns N_{HC} (lower left), mean bandedness α (lower right), characterizing OD layouts during development. At $t = 4\tau$ OD columns have segregated. After 10τ , columns start reorganizing and merge progressively towards a stationary stripe-like layout. Whereas Λ , and N_{HC} reach near mature levels around 10τ the bandedness α increases during the entire time course, capturing the slow reorganization of OD columns.

Video 2. Zigzag reorganization in the EN model after instantaneous isotropic area increase. The video displays OD maps (upper left) as well as time courses of mean column spacing Λ (upper right), number of hypercolumns N_{HC} (lower left), mean bandedness α (lower right). After instantaneous area increase at $t = 10\tau$, OD stripes bend sinusoidally, a behavior typical for ZZ reorganization. A jump in Λ is observed shortly after $t = 10\tau$. Subsequently, however, Λ decreases to almost the initial value and consequently, N_{HC} increases. α decreases, reflecting the bending of OD domains. When increasing the system size, the EN model displays all features typical for ZZ reorganization.

Video 3. OD development in the EN model with growth at near realistic growth rate (156% total area increase between $t = 0$ and $t = 100\tau$, $\eta = 0.025$, $r = 0.16$). The video displays OD maps (upper left) as well as time courses of mean column spacing Λ (upper right), number of hypercolumns N_{HC} (lower left), mean bandedness α (lower right). Despite the large area increase, column spacing Λ increases only mildly and transiently, implying a strong increase in N_{HC} . Note, that growth-induced reorganization of OD layouts shows aspects of the ZZ reorganization that is captured by a drop in α .